

## Microbiological Study of Oral Cavity of Patients Seeking Dental Treatments in Dhaka Metropolis, Bangladesh

Jannatul Ferdous Mitaa<sup>1</sup>, Tasnia Ahmed<sup>1\*</sup> 

Department of Microbiology, Stamford University Bangladesh, 51 Siddeswari Road, Dhaka 1217, Bangladesh.



### Article Information

#### Article Subject:

Oral Microbiology


 10.30699/ijmm.13.4.284

#### Corresponding author:

##### Tasnia Ahmed

Department of Microbiology,  
Stamford University Bangladesh,  
51 Siddeswari Road, Dhaka 1217,  
Bangladesh

##### Email:

[tasnia2009@yahoo.com](mailto:tasnia2009@yahoo.com)Use your device to scan  
and read the article online

### Abstract

**Background and Aims:** Periodontal diseases are very common among the people of all ages and also a proof of oral hygiene maintenance practice. A variety of microorganisms are found in human oral cavity which can often cause periodontitis, cavities, gingivitis, mineralization of tooth and other oral diseases. Current study was carried out to determine which microbes are responsible for dental caries among the patients in Dhaka along with control with no dental disease.

**Materials and Methods:** After observing the symptoms of dental caries like tooth pain, bleeding gum, bleeding while eating and brushing, weakening of teeth, mobility of teeth, swelling of gum etc.; dental plaque samples were collected and diluted in thioglycollate broth. The diluted sample was then transferred to a series of specific media including *Mitis salivarius* Agar (MSA), BD LBS Agar, *Actinomyces* Isolation Agar, NOS Spirochete medium, *P. GING* (*Porphyromonas gingivalis* Agar) and VCAT and incubated at 37°C to isolate *Streptococcus* spp., *Lactobacillus* spp., *Actinomyces* spp., *Treponema* spp. and *Capnocytophaga* spp. One sample was taken from a healthy patient as a control test harboring no pathogenic microbial growth.

**Results:** According to current study, the most common microbes which can initiate these dental diseases include *Streptococcus mutans*, *Lactobacillus* spp., *Capnocytophaga* spp., *Porphyromonas gingivitis*, *Actinomyces* spp. etc. After detecting the mentioned microorganisms, different procedures were prescribed for the patients including scaling, root canal, surgical removal of teeth (tooth extraction) etc.

**Conclusion:** Oral hygiene should be maintained not only by the patients, but also by every person to lessen the risk of dental disease as well as the early detection of it to prevent the loss of tooth.

**Keywords:** Biofilms; Caries; Dental plaque; Periodontitis

Received: 2019/08/09

Accepted: 2019/10/02

Available online: 2019/10/02

Copyright © 2019. This is an open-access article distributed under the terms of the Creative Commons Attribution- 4.0 International License which permits Share, copy and redistribution of the material in any medium or format or adapt, remix, transform, and build upon the material for any purpose, even commercially.

### How to cite this article:

Ferdus Mitaa J, Ahmed T. Microbiological study of oral cavity of patients seeking dental treatments in Dhaka Metropolis, Bangladesh. Iran J Med Microbiol. 2019; 13 (4) :284-293

Download citation: [BibTeX](#) | [RIS](#) | [EndNote](#) | [Medlars](#) | [ProCite](#) | [Reference Manager](#) | [RefWorks](#)Send citation to:  [Mendeley](#) |  [Zotero](#) |  [RefWorks](#)

## Introduction

Human body harbors different types of microorganisms located at different sites. In some areas there prevails a complex mixture of microorganisms which have inter-connections and function together. The functional characteristics of the complex of microbes often differ from their individual characteristics. In human body one such area harboring a complex mixture of microorganisms is oral cavity. Nearly 700 types of microorganisms have been found to be present in the oral microbe complex ecosystem which mainly takes part in dental diseases mostly like dental caries, periodontitis etc. which are very common infections in human seeking dental treatments (1-6). Some microorganisms of oral cavity can also cause systemic diseases like pneumonia, endocarditis, osteomyelitis etc. if they disseminate via the blood stream to distal body parts (7,8). Dental diseases like tooth decay has been prevailing from prehistoric ages which drastically increased after the increased consumption of refined sugar (9). W. D. Miller was the first pioneer who proposed the presence of oral organisms or germs in oral cavity of human who can ferment carbohydrates into acid which is responsible for tooth decay. *Streptococcus mutans* was the first bacteria which was isolated by J. K. Clarke (10,11). In different studies it was found that *Actinomyces* species can cause dental diseases (12). Dental plaque occurs due to the rapid proliferation of dental microorganisms producing biofilms which was first seen by Sir Antony Van Leuwenhoek under the microscope (13). It has been revealed that the microbes in dental plaque are responsible for the initiation of dental decays by producing acidic byproducts. Some common species found to be present in dental plaque includes *Streptococcus* spp., *Actinobacillus* spp., *Actinomyces* spp., *Porphyromonas* spp., *Treponema* spp. etc. (14). Dental plaque is formed by the actions of biofilms which is a mass of bacterial community where different microbiota can inhabit together in a gelatinous matrix. Because of the gelatinous material, the microbes are resistant to external hazards. There is also extensive metabolites exchanges, complex interactions among themselves which is unique for biofilms and cannot be seen when these species are living alone (15).

Supragingival plaque has been showed to be dominated mostly by Gram-positive bacteria like *Streptococcus sanguinis*, *Streptococcus mutans*, *Streptococcus salivarius*, *Streptococcus mitis* and lactobacilli. Subgingival plaque is harbored by gram-negative anaerobic bacteria like *Aggregatibacter (Actinobacillus) actinomycetemcomitans*, *Tannerella forsythia*, *Fusobacterium nucleatum*, *Eikenella corrodens*, *Campylobacter* spp., *Capnocytophaga* spp., *Porphyromonas gingivalis*, *Prevotella intermedia*, and oral spirochetes like *Treponema denticola* (16). The treatments for dental diseases generally include surgical approaches like root canal, filling, surgical removal of teeth and so on. But at the same time it is important to detect the cariogenic bacteria which damage and demineralize the tooth by producing acids and enhancing the ways of re-mineralization of tooth. A new approach to decrease the dental problem is to introduce probiotics in mouth instead of killing all microbes from oral cavity. Because oral cavity includes a wide variety of microbial population, killing some microbes may not lessen the problem, because other oral microbial flora can replace in the new vacant places. But if we use probiotics which don't initiate plaque formation and also can produce bacteriocin type products which can kill other plaque forming bacteria can be more useful in maintaining healthy oral cavity (17,18). In our current study we randomly selected 9 patients who came to seek dental assistance. The symptoms were first enlisted and microbiological analysis was performed to determine the presence of microorganisms in the specific infected area of oral cavity and then necessary treatment was suggested.

## Materials and Methods

### Sample collection and processing

Firstly, the physiological conditions (symptoms associated with dental diseases) for which patients were seeking dental check up along with the specific clinical symptoms identified by the healthcare personnel were enlisted. For microbiological analysis, dental plaque samples from twelve patients were collected in September, 2018 after asking their consent in a local dental

health care center in Mirpur, Dhaka, Bangladesh. Among them one sample was from healthy individual as control. There was no ethical code for the research and no further involvement of any organization regarding the ethical code was not necessary. Sampling was done by using sterile forceps or toothpicks and transferred into 2 mL of reduced transport fluid medium (0.4% agar, 0.15% thioglycollate / phosphate buffered saline and stored at 4°C until plating onto specific agar plates. While studying the sample it was transferred to 9 mL thioglycollate broth and incubated for 24 hours and then serially diluted for inoculation into specific plates for identification of several dental plaque forming bacterial isolates in case of spreading plate technique. And for streaking, the original sample that was collected was used to directly inoculate onto the agar plates (19,20).

#### Isolation of *Streptococcus* spp.

For isolation of *Streptococcus* spp., Mitis Salivarius Agar (MSA) was used. This media is especially used for isolating *Streptococcus mitis*, *Streptococcus salivarius* from mixed cultures found in dental plaque samples. After incubation period at 37°C for 18-48 hours, presence of specific microorganisms was observed. *Streptococcus mutans* produce undule-shaped colonies, with frosted-glass appearance due to producing dextran from sugar. *Streptococcus mitis* could be identified as light blue colored, small and flat colonies. And *Streptococcus salivarius* could be identified as sticky, mucoid, gum-drop like colonies.

#### Isolation of *Lactobacillus* species

BD LBS Agar (*Lactobacillus* identification agar) is used for the detection of *Lactobacillus* spp. This media is also known as Rogosa Agar. Collected dental samples were introduced directly onto this agar followed by four quadrant streaking method to determine the presence of *Lactobacillus* spp. The dental samples were collected by dental floss can also be inoculated by spreading 0.1 mL over the media. Inoculated plates were incubated at 37°C for 48 to 72 hours in anaerobic condition. After incubation, *Lactobacillus* spp. could be determined

by observing medium to large sized, white colonies onto the media.

#### Isolation of *Actinomycetes*

Collected dental floss sample (0.1 mL) was inoculated onto *Actinomycetes* Isolation Agar and incubated at 37°C for 3 to 7 days because they are slow growing microbes. After incubation *Actinomycetes* spp. can be identified as mucoid, fungi like filamentous growth characteristics onto the agar plates.

#### Isolation of *Spirochetes*

For isolating *Treponema* species like *Treponema amylovorum*, *Treponema denticola*, *Treponema maltophilum*, *Treponema medium*, *Treponema pectinovorum*, *Treponema socranskii*, and *Treponema vincentii* etc. which can be found in dental plaques are cultured in NOS Spirochete medium. After inoculation onto the NOS medium by streaking, the plates were incubated at 37°C anaerobically for 7 days. In current study we inoculated dental plaque sample onto the NOS agar medium and the Spirochete grew in NOS agar as a white, hazy, cottony growth.

#### Isolation of *Porphyromonas gingivalis*

*P. GING* media has been used to isolate *Porphyromona gingivalis* from dental plaque samples. The media was inoculated directly with the broth where the dental plaque sample was directly suspended and then streak plate method was applied. The plates were then placed in an anaerobic atmosphere and incubated at 37°C for 48 hours.

#### Isolation of *Capnocytophaga* species


VCAT media was used in this study to isolate *Capnocytophaga* spp. from dental plaques. Samples were streaked onto VCAT agar and incubated anaerobically at 37°C for 48 hours. Colonies are convex or flat and often slightly yellow, show regular or spreading edges.

Table 1- Symptoms and microorganisms responsible for dental plaque formation among patients.


Patient No.	Clinical Symptoms	<i>Actinomyces</i>	<i>Lactobacillus</i> spp.	<i>Capnocytophaga</i> spp.	<i>Streptococcus mutans</i>	<i>Treponema denticola</i>	<i>Porphyromonus gingivalis</i>	Treatment suggested
01	<ul style="list-style-type: none"> <li>Severe Pain at 6</li> <li>It hampered sleep</li> <li>Pain reduced for some hour and again starts.</li> <li>Per apical X-ray showed pulpitis with per apical lesion in left 6 tooth.</li> </ul>	-	++	-	++	-	-	Left 6 Root canal treatment performed followed by porcelain crown. Scaling and oral hygiene instruction done. Suggested for check up after 3 months.
02	<ul style="list-style-type: none"> <li>Severe Pain at 6</li> <li>It hamper his sleep</li> <li>Pain reduce for some hour and again starts.</li> <li>Per apical x-ray showed pulpitis with per apical lesion in left 7 no tooth.</li> </ul>	-	++	-	++	-	-	Left 6 Root canal treatment performed followed by porcelain crown.
03	<ul style="list-style-type: none"> <li>Severe Pain at 6</li> <li>It hamper his sleep</li> <li>Pain reduce for some hour and again starts.</li> <li>Per apical x-ray showed pulpitis with per apical lesion in left 4 and 5 no tooth.</li> </ul>	-	++	-	++	-	-	Left 4,5 Root canal treatment performed followed by porcelain crown. Scaling and oral hygiene instruction done. Suggested for check up after 3 months.
04	<ul style="list-style-type: none"> <li>Severe Pain upper right 7 no tooth</li> <li>It hamper his sleep</li> <li>Pain reduce for some hour and again starts.</li> <li>Per apical x-ray showed pulpitis with per apical lesion in upper right 7 no tooth.</li> </ul>	-	++	-	++	-	-	Right Upper 7 Root canal treatment performed followed by porcelain crown. Scaling and oral hygiene instruction done. Suggested for check up after 3 months.
05	<ul style="list-style-type: none"> <li>pain at left 3,4,5,7, and right 2,6</li> <li>second degree mobility</li> <li>Peripheral X-ray showed horizontal and alveolar bone loss.</li> </ul>	-	-	-	-	++	++	Scaling done. Systemic abnormalities should be treated and oral hygiene instruction done. Suggested for check up after 3 months.
06	<ul style="list-style-type: none"> <li>pain at left 1,2,3 and right 1,2,3</li> <li>second degree mobility</li> <li>Peripheral X-ray showed horizontal and alveolar bone loss.</li> </ul>	-	-	-	-	++	++	Scaling done. Systemic abnormalities should be treated and oral hygiene instruction done. Suggested for check up after 3 months.

Patient No.	Clinical Symptoms	<i>Actinomyces</i>	<i>Lactobacillus</i> spp.	<i>Capnocytophaga</i> spp.	<i>Streptococcus mutans</i>	<i>Treponema denticola</i>	<i>Porphyromonas gingivalis</i>	Treatment suggested
07	<ul style="list-style-type: none"> <li>• Bad smell</li> <li>• Bleeding Gum</li> <li>• Bleeding Continue While Eating and brushing</li> <li>• Gingival swelling</li> </ul>	++	-	++	-	-	-	Scaling, Oral hygiene instruction, Review after 6 months.
08	<ul style="list-style-type: none"> <li>• Bad smell</li> <li>• Bleeding Gum</li> <li>• Bleeding Continue While Eating and brushing</li> <li>• Gingival swelling</li> </ul>	++	-	++	-	-	-	Scaling, Oral hygiene instruction, Review after 6 months.
09	<ul style="list-style-type: none"> <li>• pain at left,3 and right 2</li> <li>• second degree mobility</li> <li>• Peripheral X-ray showed horizontal and alveolar bone loss.</li> </ul>	-	-	-	-	++	++	Scaling done. Systemic abnormalities should be treated and oral hygiene instruction done. Suggested for check up after 3 months.
10	<ul style="list-style-type: none"> <li>• Severe Pain at 7</li> <li>• It hamper his sleep</li> <li>• Pain reduce for some hour and again starts.</li> <li>• Per apical x-ray showed pulpitis with per apical lesion in left 7 no tooth.</li> </ul>	-	-	-	++	-	-	Left 7 Root canal treatment performed followed by porcelain crown.
11	<ul style="list-style-type: none"> <li>• Bad smell</li> <li>• Bleeding Gum</li> <li>• Bleeding Continue While Eating and brushing</li> <li>• Gingival swelling</li> </ul>	++	-	++	-	-	-	Scaling, Oral hygiene instruction, Review after 6 months.
12	<ul style="list-style-type: none"> <li>• No caries, No bleeding gum</li> <li>• Came for dental checkup</li> </ul>	-	-	-	-	-	-	No treatments needed.







**Figure 1.** Adult human as 32 teeth having distinct names for them. As much as their position is situated in the corner of mouth where it is difficult to reach for brushing, the risk for developing infections also increases simultaneously. Most infections occur Premolar and molar teeth.



**Figure 2.** Different parts of a tooth has been indicated in the figure. To determine the disease conditions of a tooth, it is necessary to identify separate parts of a tooth because infection at these parts can be caused by separate pathogenic bacteria. The resulting infection in different portions might need distinct dental procedures to correct the situation.



**Figure 3.** The dental caries is initiated by different ways. Different parts of the tooth can be affected during the progression of dental caries. As the condition becomes worse by further dissemination of the caries, the infection and disease symptoms increase. First the dental caries is initiated by the biofilm of pathogenic bacteria thriving in oral cavity if adequate cleaning is not regularly maintained. This initiation can be detected as a small caries on the enamel. If the condition is not corrected by proper treatment, the caries further progresses inside the tooth (dentin and then pulp). During this stage the microorganisms reach the pulp and abscess is formed followed by inflammation and deepened pocket calculus.



**Figure 4.** It is always better to take some preventive measures to decrease the chance of dental plaque formation. If the consumed food contains lots of sugar content it is very important to brush teeth as soon as possible to stop the accumulation of microbes on the tooth surface. Oral hygiene should be maintained strictly. Teeth should be brushed twice a day and after eating sweet food. Yearly dental check up is recommended to make early detection of caries and providing early treatment before making it worse.

## Results

Microbial biofilm formation is very common phenomenon in teeth which leads to the dental diseases starting with dental caries. Molar teeth (Figure 1) are most prone to such problems due to the difficulties in cleaning different food particles. Different parts of tooth are damaged during the progression of dental diseases (Figures 2, 3). After observing the disease symptoms as well as the microbiological analysis of the dental plaques, it has been revealed that certain types of microorganisms together forming dental plaques are responsible for certain symptoms (Table 1). *Lactobacillus* spp. and *Streptococcus mutans* were present in sample no. 1, 2, 3, 4 and in all these four cases there prevailed same types of symptoms including severe pain, restless night due to intensive pain. In all cases pain reduced for several hours and again started. *Lactobacillus* spp. and *Streptococcus mutans* together formed dental plaques in different tooth in these patients and maintaining oral hygiene was suggested. Root canal for the infected tooth was prescribed to remove the deep plaques and porcelain crown can be used depending on the condition of the tooth. *Actinomyces* spp. was present in sample no. 7, 8 and 11. *Capnocytophaga* spp. (sample 7, 8), *Treponema denticola* and *Porphyromonas gingivalis* (sample 5,6,9) were also present. Control sample (healthy individual) was showed no growth of such bacteria.

## Discussion

*Lactobacillus* spp. is responsible for root caries in adults mostly. They can attach to the tooth surface easily and the most importantly they can co-aggregate with other species of microorganisms (especially *Streptococcus* spp.) and can initiate the biofilm production where prevails a complex interrelated metabolic reactions and functions (21). *Streptococcus mutans* in tooth surface utilizes sugar to produce lactic acid using the enzyme glucanucrase. The produced acid causes the mineralization of teeth enamel initiating the dental caries (22).

In patient no 5, there was pain at left 3,4,5,7, and right 2,6 teeth due to dental plaque formation

and the microbiological analysis revealed the presence of *Treponema denticola* and *Porphyromonas gingivalis*. Dental scaling was suggested to remove the plaques and further maintenance of hygiene was advised. In patient no 5, there was also pain at left 1,2,3 and right 1,2,3 and the plaque forming microorganisms were *Treponema denticola* and *Porphyromonas gingivalis* like patient no 5. *Treponema denticola* is responsible for gum inflammation and periodontal disease. They can survive extreme harsh environment in mouth and make biofilms. In biofilms, they generally co-aggregate with *Porphyromonas gingivalis* (23,24). *Porphyromonas gingivalis* is one of the most common pathogen responsible for periodontic diseases. 85.75% samples of dental plaque showed the presence of this bacteria. It can survive in deep cavities where sugar availability is low and they can utilize amino acids instead of sugar. They can communicate with other bacteria increasing the advancement of the disease (25).

Patients 7 & 8 both showed similar symptoms like bad smell, bleeding Gum, bleeding continued while eating and brushing. Gingival swelling was also observed. In both patients, responsible microorganisms forming dental plaque were *Actinomyces* spp. and *Capnocytophaga* spp. *Capnocytophaga* strains are isolated from dental plaques with other bacterial periodontal species of bacteria. This condition is responsible for alveolar bone loss, attachment loss, tooth mobility and tooth loss (26). *Actinomyces* spp. can produce black stain or tooth discoloration, prone to calcification. They can also cause dental abscess (Figure 3).

Several studies were carried out before where similar microorganisms were identified causing distinct periodontal and gum disease (27-30).

Treatments include dental filling using amalgam which is silver-gray colored material. For better visualization, tooth colored filling components are also available. If the caries is big enough to be filled properly, artificial crown or cap is used as a cover over the damaged tooth. When the infection reaches the deepest part of tooth infecting the



nerve n pulp, root canal is performed to remove the infected portion and filled with sealing material. And then a crown is placed over it, if necessary.

## Conclusion

Dental caries is a very common dental problem occurring in the children as well as in adults. Many different types of microorganisms are responsible for dental caries. *Streptococcus mutans*, *Lactobacillus* spp., *Actinomyces* spp., *Capnocytophaga* spp., *Porphyromonas* spp. etc bacteria have been identified from the patients subjected to microbiological analysis in this study. They can co-aggregate with each other enabling themselves to become more virulent and increase the disease symptoms and dental damages as well. Scaling, root canal, fill ups and finally surgical removal of damaged tooth are the main treatments for the dental problems. Oral hygiene maintenance can decrease such problems.

## Acknowledgements

Authors are thankful to the researchers of the related studies which have been cited in the text.

## Conflict of Interest

The authors reported no conflict of interest.

## Consent

Patients' consent was taken before sampling of the dental plaques by the dentist while working in the chamber.

## References

1. Aas JA, Paster BJ, Stokes LN, Olsen I, Dewhirst FE. Defining the Normal Bacterial Flora of the Oral Cavity. *J. Clin. Microbiol* 2005 43:5721-5732. [[DOI:10.1128/JCM.43.11.5721-5732.2005](https://doi.org/10.1128/JCM.43.11.5721-5732.2005)] [[PMID](#)] [[PMCID](#)]
2. Nishihara T, Koseki T. Microbial etiology of periodontitis. *Periodontology* 2000 2004 36: 14-26. [[DOI:10.1111/j.1600-0757.2004.03671.x](https://doi.org/10.1111/j.1600-0757.2004.03671.x)] [[PMID](#)]
3. Kesic L, Milasin J, Igetic M, Obrandovic R. Microbial etiology of periodontal disease- mini review. *Medicine and Biology* 2008 15(1): 1-6.
4. Paster BJ, Boches SK, Galvin JL, Ericson RE, Lau CN, Levanos VA, Sahasrabudhe A, Dewhirst FE. Bacterial Diversity in Human Subgingival Plaque. *J. Bacteriol* 2001 183:3770-3783. [[DOI:10.1128/JB.183.12.3770-3783.2001](https://doi.org/10.1128/JB.183.12.3770-3783.2001)] [[PMID](#)] [[PMCID](#)]
5. Paster BJ, Olsen I, Aas JA, Dewhirst FE. The breadth of bacteria diversity in the human periodontal pocket and other oral sites. *Periodontology* 2000 2006 42:80-87. [[DOI:10.1111/j.1600-0757.2006.00174.x](https://doi.org/10.1111/j.1600-0757.2006.00174.x)] [[PMID](#)]
6. Goncalves C, Soares GMS, Faveri M, Perez-Chaparro PJ, Lobao E, Figueiredo LC, Baccelli GT, and Feres M. Association of three putative periodontal pathogens with chronic periodontitis in Brazilian subjects. *Journal of Applied Oral Science* 2016 24(2): 181-185. [[DOI:10.1590/1678-775720150445](https://doi.org/10.1590/1678-775720150445)] [[PMID](#)] [[PMCID](#)]
7. Berbari EF, Cockerill III FR, and Steckelberg JM. Infective endocarditis due to unusual or fastidious microorganisms. *Mayo Clin. Proc* 1997 72:532-542. [[DOI:10.4065/72.6.532](https://doi.org/10.4065/72.6.532)] [[PMID](#)]
8. Gomes-Filho IS, Passos JS, Seixas da Cruz S. Respiratory disease and the role of oral bacteria. *Journal of Oral Microbiology* 2010 2:10.3402/jom.v2i0.5811. [[DOI:10.3402/jom.v2i0.5811](https://doi.org/10.3402/jom.v2i0.5811)] [[PMID](#)] [[PMCID](#)]
9. Bifulco M, Amato M, Gangemi G, Marasco M, Caggiano M, Amato A, Psanti S. Dental care and dentistry practice in the Medieval Medical School of Salerno. *BDJ* 2016 221: 87-89. [[DOI:10.1038/sj.bdj.2016.528](https://doi.org/10.1038/sj.bdj.2016.528)] [[PMID](#)]
10. Miller WD. *Graphische Anstalt Schuler AG. Biel, Switzerland: 1890. The micro-organisms of the human mouth.*
11. Xiao C, Ran S, Huang Z, Liang J. Bacterial Diversity and Community Structure of Supragingival Plaques in Adults with Dental Health or Caries Revealed by 16S Pyrosequencing. *Frontiers in Microbiology* 2016 7:1145. [[DOI:10.3389/fmicb.2016.01145](https://doi.org/10.3389/fmicb.2016.01145)]
12. Könönen E, Wade WG. Actinomyces and related organisms in human infections. *Clin Microbiol Rev* 2015 28(2): 420-442. [[DOI:10.1128/CMR.00100-14](https://doi.org/10.1128/CMR.00100-14)] [[PMID](#)] [[PMCID](#)]
13. Gest H. The discovery of microorganisms by Robert Hooke and Antoni van Leeuwenhoek, Fellows of The Royal Society. *Notes and Records*

- of the Royal Society of London 2004 58:187-201. [DOI:10.1098/rsnr.2004.0055] [PMID]
14. Kolenbrander PE. Oral microbial communities: Biofilms, Interactions, and Genetic Systems. *Annu. Rev. Microbiol* 2000 54:413-437. [DOI:10.1146/annurev.micro.54.1.413] [PMID]
  15. Lappin Scott H, Burton S, Stoodley P. Revealing a world of biofilms - the pioneering research of Bill Costerton. *Nature Reviews Microbiology* 2014 12(11): 781-787. [DOI:10.1038/nrmicro3343] [PMID]
  16. He X, Shi W. Oral Microbiology: Past, Present and Future. *International Journal of Oral Science* 2009 1(2): 47-58. [DOI:10.4248/ijos.09029] [PMID] [PMCID]
  17. Meurman JH, Stamatova I. Probiotics: contributions to oral health. *Oral Dis* 2007 13:443-451. [DOI:10.1111/j.1601-0825.2007.01386.x] [PMID]
  18. Guarner F, Perdigon G, Coerthier S, Salminen B, Morelli L. Should yoghurt cultures be considered probiotic? *Br. J. Nutr* 2005 93:783-786. [DOI:10.1079/BJN20051428] [PMID]
  19. Chandrabhan D, Hemlata R, Renu B, Pradeep V. Isolation of dental caries bacteria from dental plaque and effect of tooth pastes on acidogenic bacteria. *Open Journal of Medical Microbiology* 2012 2: 65-69. [DOI:10.4236/ojmm.2012.23009]
  20. Salam MA, Senpuku H, Nomura Y, Matin K, Miyazaki H and Hanada N. "Isolation of Opportunistic Pathogens in Dental Plaque, Saliva and Tonsil Sample from Elderly," *Japanese Journal of Infectious Diseases* 2001 54(5): 193-195.
  21. Badet C, Thebaud NB. Ecology of Lactobacilli in the oral cavity. *Open Microbiol J* 2008 2: 38-48. [DOI:10.2174/1874285800802010038] [PMID] [PMCID]
  22. Loesche WJ. Ch. 99: Microbiology of Dental Decay and Periodontal Disease. In Baron S; et al. *Baron's Medical Microbiology* (4th ed.). University of Texas Medical Branch. 1996.
  23. Kuramitsu HK, Chen W, Ikegami A. Biofilm formation by the periodontopathic bacteria *Treponema denticola* and *Porphyromonas gingivalis*. *J Periodontol* 2005 76(11): 2047-51. [DOI:10.1902/jop.2005.76.11-S.2047]
  24. Catherine A. Brissette1, Sheila A. Lukehart. "Mechanisms of Decreased Susceptibility to  $\beta$ -Defensins by *Treponema denticola*. Infection and Immunity 2007 75(5): 2307-2315. [DOI:10.1128/IAI.01718-06] [PMID] [PMCID]
  25. Bostanci N, Belibasakis GN. *Porphyromonas gingivalis*: an invasive and evasive opportunistic oral pathogen. *FEMS Microbiol. Lett* 2012 333(1) 1-9. [DOI:10.1111/j.1574-6968.2012.02579.x] [PMID]
  26. Loannou AL, Kotsakis GA, Hinrichs JE. *World J Clin Cases* 2014 2(12): 822-827. [DOI:10.12998/wjcc.v2.i12.822] [PMID] [PMCID]
  27. Slots J, Moenbo D, Langebaek J, Frandsen A. Microbiota of gingivitis in man. *Scandinavian Journal of Dental Research*. 1978 174-181. [DOI:10.1111/j.1600-0722.1978.tb01929.x] [PMID]
  28. Moore WE, Holdeman LV, Smibert RM, Hash DE, Burmeister JA, Ranney RR. Bacteriology of severe periodontitis in young adult humans. *Infection and immunity*. 1982 38:1137-1148. [DOI:10.1128/IAI.38.3.1137-1148.1982] [PMID] [PMCID]
  29. Popova C, Panova DV, Panov V. Microbiology of periodontal diseases. A review. *Biotechnology & Biotechnological Equipment*. 2013 27:3754-3759. [DOI:10.5504/BBEQ.2013.0027]
  30. Darout Al. Oral bacterial interaction in periodontal health and disease. *Journal of Dentistry and Oral Hygiene*. 2014 6:51-57. [DOI:10.5897/JDOH2014.0127]