

Bifidobacterium bifidum Association with Changes in miR-196a/b-5p and Its Target Genes in HT-29 Colorectal Cancer Cells

Arghavan Sabzevari¹ , Somayeh Farahmand^{1*} , Mohammad Amin Nicknezhad² 

1. Department of Biology, Payame Noor University (PNU), Tehran, Iran
2. Department of Microbiology, Islamic Azad University, Shahrekord Branch, Shahrekord, Iran

ABSTRACT

Background and Aim: Probiotics, particularly species of *Bifidobacterium*, have shown potential anticancer properties through mechanisms involving immune modulation, regulation of apoptosis, and production of bioactive metabolites. This study aimed to examine the cytotoxic effects of post-fermentation medium (PFM) and cell extract (CE) derived from *Bifidobacterium (B.) bifidum* on colorectal cancer (CRC) cells and to evaluate their impact on the expression of cancer-related microRNAs (miRNAs) and target genes.

Materials and Methods: HT-29 colorectal cancer cells were exposed to different concentrations (0–50% v/v) of PFM and CE. Cytotoxicity was assessed using the MTT assay. The expression levels of miR-196a-5p and miR-196b-5p and their target genes (*HOXB8*, *IGF2BP3*, and *E2F7*) were quantified using RT-qPCR.

Results: PFM induced a dose-dependent reduction in HT-29 cell viability, with an IC₅₀ of approximately 35%, while CE displayed minimal cytotoxicity. PFM also upregulated miR-196a-5p and miR-196b-5p, particularly at 35% and 50%, whereas CE had no significant effect. Correspondingly, PFM was associated with significant downregulation expression of *HOXB8* and *IGF2BP3*. No meaningful changes were observed in *E2F7* expression or in CE-treated cells.

Conclusion: *Bifidobacterium bifidum*-derived PFM demonstrated anticancer activity by reducing cancer cell viability and showing an association with changes in miRNA and target gene expression. However, these findings are correlative and do not establish a causal, miRNA-mediated mechanism. Further functional studies (e.g., using miRNA mimics/inhibitors or gene knockdown) are required to determine whether the observed gene expression changes are directly mediated by miRNA modulation, and *in vivo* validation is also warranted.

Keywords: *Bifidobacterium bifidum*, Colorectal Neoplasms, Fermentation Products, Gene Expression Regulation, MicroRNAs

Received: 2025/10/13;

Accepted: 2026/01/17;

Published Online: 2026/02/28;

Corresponding Information: Somayeh Farahmand*, Department of Biology, Payame Noor University (PNU), Tehran, Iran & Email: s.farahmand@pnu.ac.ir



Copyright © 2026. This is an original open-access article distributed under the terms of the [Creative Commons Attribution-noncommercial 4.0 International License](#) which permits copy and redistribution of the material just in noncommercial usages with proper citation.

Use a device to scan and read the article online



Sabzevari A, Farahmand S, Niknezhad M A. *Bifidobacterium bifidum* Association with Changes in miR-196a/b-5p and Its Target Genes in HT-29 Colorectal Cancer Cells. Iran J Med Microbiol. 2026;20 (1):1-11.

Download Citation:

[RIS](#) | [EndNote](#) | [Mendeley](#) | [BibTeX](#) |

1. Introduction

Colorectal cancer (CRC) is one of the leading causes of cancer-related morbidity and mortality worldwide. This malignancy arises from a complex interaction between genetic susceptibility, environmental influences, and disturbances in the gut microbiota (1). Recent developments in cancer research have emphasized

the crucial importance of the intestinal microbiome in modulating physiological processes relevant to tumor development. Among the various microbial inhabitants *Bifidobacterium (B.) bifidum*, a prominent probiotic species in the human gut, has gained increasing attention for its potential involvement in cancer-related mechanisms. *B. bifidum*, widely

recognized for its beneficial effects on gastrointestinal health, exhibits several properties that may contribute to its anticancer activity, including strengthening the intestinal barrier, modulating host immune responses, and maintaining microbial homeostasis (2-4).

Notably, emerging evidence suggests that the influence of probiotics extends to the regulation of gene expression through epigenetic mechanisms, particularly via the modulation of microRNAs (miRNAs) (5).

The miRNAs are tiny non-coding RNAs that influence the regulation of genes expression at post-transcriptional level. This regulation plays critical roles in various health and disease states. Among them, miR-196a/b-5p has attracted noteworthy consideration due to its oncogenic role in several types of cancer. Increased levels of miR-196a/b-5p have been related to enhanced tumor cell growth, invasion, and resistance to programmed cell death (6, 7). In CRC, aberrant levels of miR-196a/b-5p contribute to the maintenance of an oncogenic phenotype (8, 9). Thus, targeting dysregulated miRNA expression may provide new therapeutic strategies for the colorectal cancer.

Several target genes of miR-196a/b-5p have been implicated in tumorigenesis. *HOXB8*, a homeobox transcription factor, is overexpressed in CRC and enhances proliferation of cells and their invasive behavior metastasis, and chemotherapy resistance via the Wnt/ β -catenin and STAT3 signaling pathways. Insuline-like growth factor 2 mRNA-binding protein 3 (IGF2BP3) stabilizes oncogenic mRNAs by recognizing their N⁶-methyladenosine (m⁶A) modifications and is related to tumor growth, metastasis, and unfavorable outcomes across multiple cancers (10). The *E2F7* gene, a transcriptional regulator of the E2F family, is frequently upregulated in CRC, where it modulates cell cycle progression and may influence proliferation, although its precise role remains context-dependent (11). Understanding the modulation of these genes may clarify the mechanisms by which probiotics influence cancer biology.

This study investigates the regulatory effects of *B. bifidum* on miR-196a/b-5p in HT-29 CRC cells. It also examines how these miRNAs influence the level of important target genes expression that are involved in tumor progression.

The HT-29 cell line is a well-established *ex vivo* model for studying colorectal tumor biology due to its tumorigenic characteristics (12, 13). Despite increasing interest in probiotic-cancer interactions, the molecular mechanisms by which *B. bifidum* modulates miRNAs—particularly miR-196a/b-5p and

its target genes remain largely unexplored. This study represents the first inquiry into the impact of *B. bifidum* on the level expression of miR-196 a/b-5p and its key target genes—*HOXB8*, *E2F7*, and *IGF2BP3*—which are critically involved in CRC cell proliferation, invasion, and cell death. Considering the essential role of miRNAs in cancer biology and the growing interest in microbiota-based therapeutics, elucidating how *B. bifidum* modulates miR-196a/b-5p and its downstream targets could pave the way for novel miRNA-targeted therapies or improve existing CRC treatments.

2. Materials and Methods

2.1 Bacterial Strain and Culture Conditions

Bifidobacterium (B.) bifidum (strain 1644 PTCC) was obtained in a pure, lyophilized form from the Iranian Industrial Microbial Culture Collection (Iran Scientific and Industrial Research Organization). The bacterium was revived and grown in de Man, Rogosa, and Sharpe (MRS) broth (Merck, Germany). The cultures were incubated anaerobically at 37°C for 72 hr using an anaerobic jar and gas pack system (AnaeroGen™, Oxoid, UK).

2.2 Preparation of Post-Fermentation Medium (PFM)

To prepare the post-fermentation medium (PFM), an active culture of *B. bifidum* was inoculated into fresh MRS broth and incubated for 24 hr at 37°C under anaerobic conditions. The culture was centrifuged at 10,000×g for 15 min at 4°C. In the next step, the cell-free supernatant was obtained. The pH level was calibrated to 7.0 ± 0.1 using 1N NaOH or HCl, measured with a pH meter (Metrohm 827 pH Lab, Switzerland). Filtration through a 0.22 μm syringe filter (Sartorius, Germany) was employed to sterilize the supernatant under a Class II laminar flow cabinet. The filtered PFM was kept in sterile, capped tubes at -20°C until further use. For the cell treatment experiments, HT-29 cells treated with fresh MRS medium (without bacterial culture) were included as negative control to account for any effect of the medium alone (14, 15).

2.3 Preparation of Cell Extract (CE)

To prepare the bacterial cell extract (CE), *B. bifidum* was cultured as described above. After incubation, the cells were collected by centrifugation at 10,733×g for 15 min at 4°C. The cell pellets were washed three times with sterile phosphate-buffered saline (PBS) and centrifuged again. The washed cells were lysed using an ultrasonic bath sonicator (5 min, 50% amplitude; pulse on: 6 s; off: 2 s) keeping in ice bath. Further

homogenization was performed using a manual homogenizer to ensure complete disruption. The lysate was adjusted to pH 7.1 ± 0.1 and sterilized through 0.22 μm filtration. The final CE was stored at -20°C until utilization (15).

2.4 Cell Culture and Treatment

The HT-29 human colorectal adenocarcinoma cell line (NCBI code: C466) was obtained from the Pasteur Institute of Iran. Cells were cultured in RPMI-1640 medium (Gibco, USA) supplemented with 20% fetal bovine serum (FBS) and 1% penicillin-streptomycin (P/S), under standard conditions (37°C in a humidified incubator with 5% CO_2).

For the treatment, cells were seeded into 6-well plates (5×10^5 viable cells/well) (counted using the trypan blue exclusion method). Following 24-hr incubation to reach semi-confluency, the medium was replaced with fresh RPMI-1640 containing 10–50% (v/v) PFM or CE. Each treatment condition was performed in three independent biological replicates ($n = 3$), conducted on different days with freshly seeded HT-29 cells.

Within each biological replicate, MTT measurement was performed in technical triplicates, and qPCR assay was conducted in technical duplicates to ensure the accuracy and reproducibility (16). Cells were incubated for 48 hr under standard conditions before harvesting, and media were removed prior to further analysis. Data are presented as the mean \pm SD of the three biological replicates (15).

2.5 MTT Cell Viability Assay

To assess the cytotoxic effects of PFM and CE, MTT assay was carried out wherein cells were placed 96-well plate (10,000 cell/well) and allowed to incubate overnight. On the subsequent day, the cells received treatments with PFM or CE at concentrations between 10% and 50% (v/v) for 48 hr. Then, 20 μL of MTT solution (5 mg/mL in PBS; Sigma-Aldrich, Cat# M5655) was added to each well and incubated for 4 hr at 37°C . Following incubation, the medium was gently aspirated, and 200 μL of dimethyl sulfoxide (DMSO; Sigma-Aldrich) was added to dissolve the formazan crystals. The optical density (OD) was then measured at 570 nm, with background absorbance at 630 nm subtracted. The percentage of cell viability was

assessed with the following formula: Viability (%) = $(\text{Abs_sample} / \text{Abs_control}) \times 100$.

2.6 RNA Extraction and Real-Time qPCR

After 48 hr of treatment, total RNA was extracted using the Pars Tous RNA Extraction Kit (Cat# A101231, Iran) according to the manufacturer's instructions. The concentration and purity of the extracted RNA were evaluated spectrophotometrically.

Complementary DNA (cDNA) synthesis for mRNA targets was performed using the Ad Bio cDNA Synthesis Kit (Cat# RT-20, Korea). For miRNA analysis, cDNA was synthesized using the All-in-One miRNA First-Strand cDNA Synthesis Kit (GeneCopoeia, Cat# QP015, Canada), which includes proprietary miRNA-specific primers and a universal reverse primer. It was used strictly according to the manufacturer's protocol.

Quantitative real-time PCR (RT-qPCR) was carried out using SYBR Green Master Mix (Yekta Tajhiz Azma, Cat# YT2552, Iran) on a Rotor-Gene Q real-time PCR system (Qiagen, Germany). Primer pairs for all mRNA targets were custom-designed based on reference mRNA sequences obtained from the NCBI database. All mRNA primers were validated in our laboratory by the standard and melt curve analyses to confirm amplification efficiency (90–105%) and specificity.

GAPDH and RNU6 were used as internal reference genes for normalization of mRNA and miRNA expression levels, respectively. All qPCR reactions were performed in technical duplicates to ensure reproducibility and measurement accuracy. Relative gene expression levels were calculated using the $2^{-\Delta\Delta\text{CT}}$ method (Table 1).

2.7 Statistical Analysis

All data are presented as mean \pm standard deviation (SD) from three biological replicates ($n = 3$ independent experiments). Technical replicates (triplicates for MTT and duplicates for qPCR) were performed within each biological replicate to ensure the measurement precision. Statistical analyses were performed using GraphPad Prism 9.0. Group comparisons were evaluated by one-way ANOVA followed by Tukey's *post-hoc* test. A P -value < 0.05 was considered statistically significant. The Shapiro–Wilk test was used to assess data normality.

Table 1. Primer sequences.

Gene symbol	NCBI number	Forward primer (5'–3')	Reverse Primer (5'–3')
GAPDH	NM_0020467	CTCAGACACCATGGGAAGGTGA	ATGATCTTGAGGCTGTTGTCATA
<i>IGF2BP3</i>	NM_006547	TCGTGACCAGACACCTGATGAG	GGTGCTGCTTTACCTGAGTCAG
<i>E2F7</i>	NM_203394	TCTGAACCCGACTGTCCCTCTT	TTTGGCAGCCACATCCAGAGTG
<i>HOXB8</i>	NM_024016	GTCGCCACACAGCTCTTCCC	AATAGGAACTCCTTCTCCAGCTC
RNU6-F	NR_0030272	Proprietary (GeneCopoeia kit)	Universal primer (kit)
miR-196a-5p	MI:0000226	Proprietary (GeneCopoeia kit)	Universal primer (kit)
miR-196b-5p	MI:0001080	Proprietary (GeneCopoeia kit)	Universal primer (kit)

3. Results

3.1 Cytotoxic Effects of PFM and CE on HT-29 Cells

To assess the cytotoxic potential of post-fermentation medium (PFM) and cell extract (CE; sterile lysate of non-fermented *B. bifidum*), HT-29 cells were exposed to increasing concentrations (0–50% v/v) of each treatment for 48 hr, followed by MTT assay. HT-29 cells treated with fresh MRS medium alone were considered as negative control.

As shown in [Figure 1](#) and [Table 2](#), PFM reduced HT-29 cell viability in a concentration-dependent manner, with the most notable decline at 40% and 50% ($41\% \pm 4.2$ and $33\% \pm 5.5$, respectively; $P < 0.01$). The calculated IC_{50} was approximately 35% v/v, indicating a moderate cytotoxic threshold. In contrast, CE did not significantly affect cell viability at any tested concentration (all values $>89\%$; $P > 0.05$), suggesting that non-fermented bacterial lysates lack the cytotoxic components present in PFM.

3.2 Impact of PFM and CE on miR-196a/b-5p Expression in HT-29 Cells

To evaluate whether *B. bifidum*-derived fractions are associated with changes in miR-196a-5p and miR-196b-5p expression, HT-29 cells were exposed to different concentrations (0%, 20%, 35%, and 50% v/v) of PFM or CE for 48 hr. The expression levels were measured via RT-

qPCR and normalized to RNU6. Cells treated with fresh MRS medium alone served as negative control.

As summarized in [Figure 2](#) and [Table 3](#), PFM treatment at 35% concentration increased the expression of miR-196a-5p (1.98 ± 0.12) and miR-196b-5p (1.58 ± 0.12) significantly compared to control ($P < 0.05$). A slight decrease was observed at 50%, but expression levels remained higher than in untreated cells. CE treatment did not significantly affect miRNA expression at any tested concentration ($P > 0.05$).

3.3 Expression of miR-196a/b-5p Target Genes Following PFM and CE Treatment

To investigate the molecular mechanisms associated with observed changes in miRNA, the mRNA expression levels of three established miR-196a/b-5p targets—*HOXB8*, *IGF2BP3*, and *E2F7*—were measured in HT-29 cells exposed to various concentrations of PFM and CE for 48 hr. The expression levels were normalized to GAPDH and compared with untreated controls.

PFM treatment resulted in a significant downregulation of *HOXB8* and *IGF2BP3*, particularly at a 35% concentration, whereas *E2F7* expression remained relatively unchanged under all conditions. CE treatment did not significantly affect the expression of any target genes at any tested concentration ([Figure 3](#) and [Table 4](#)).

Table 2. The impact of CE and PFM concentrations on HT-29 cells viability.

Concentration (v/v)	PFM (%)	CE (%)
0	100	100
10	90±2	98±1.3
20	85±5	95±2
30	65±3*	96±4
40	41±4.2**	90±5
50	33±5.5**	89±2.5

Statistical significance: $P < 0.05$ (*), $P < 0.01$ (**)

Table 3. Effect of CE and PFM on expression level of miR-196a-5p and miR-196b-5p in HT-29 cells (fold change ± SD).

Concentration (v/v)	Fold change of PFM		Fold change of CE	
	miR-196b-5p	miR-196a-5p	miR-196b-5p	miR-196a-5p
0	1	1	1	1
20	1.23±0.23	1.03±0.23	1.03±0.13	1.08±0.13
35	1.98±0.12*	1.58±0.12*	1.08±0.02	1.28±0.02
50	1.79±0.31	1.6±0.31	1.19±0.21	1.25±0.21

Statistical significance: P<0.05 (*)

Table 4. Fold change in expression levels of *IGF2BP3*, *E2F7*, and *HOXB8* genes in HT-29 cells treated with CE and PFM at different concentrations (v/v).

Concentration (v/v)	Fold change of PFM			Fold change of CE		
	<i>HOXB8</i>	<i>E2F7</i>	<i>IGF2BP3</i>	<i>HOXB8</i>	<i>E2F7</i>	<i>IGF2BP3</i>
0	1	1	1	1	1	1.07
20	0.9	0.95	1.08	1.18	1.04	1.10
35	0.46**	1.11	0.77*	1.12	0.89	1.05
50	0.42	0.97	0.80	1.06	0.93	1.21

Statistical significance: P<0.05 (*), P<0.01 (**)




Figure 1. Effects of PFM and CE on HT-29 cell viability measured by MTT assay. Data are presented as mean ± SD from three independent biological replicates (n = 3). (A) PFM treatment caused a dose-dependent decrease in cell viability (IC₅₀ ≈ 35% v/v). (B) CE showed no significant effect on viability (P>0.05 vs. control). Statistical significance: P<0.05 compared to untreated control (Prepared by Authors, 2026).





Figure 2. Effects of PFM and CE on miR-196a-5p and miR-196b-5p expression in HT-29 cells. Cells were treated with 0%, 20%, 35%, and 50% (v/v) for 48 hr. Expression levels were measured by RT-qPCR and normalized to RNU6. Data are presented as fold change \pm SD from three independent biological replicates (n = 3). (A, B) PFM treatment was associated with increased miRNA expression at 35% ($P < 0.05$). (C, D) CE treatment showed no significant effect ($P > 0.05$) (Prepared by Authors, 2026).






Figure 3. Effects of PFM (A) and CE (B) on the expression of *HOXB8*, *IGF2BP3*, and *E2F7* in HT-29 cells. Gene expression was quantified by RT-qPCR and normalized to GAPDH. Data are presented as fold change \pm SD from three independent biological replicates ($n = 3$), relative to untreated control (0%). (A) PFM treatment decreased *HOXB8* and *IGF2BP3* expression at 35% and 50%, while *E2F7* expression remained unchanged. (B) CE treatment did not significantly alter gene expression at any concentration ($P > 0.05$) (Prepared by Authors, 2026).

4. Discussion

While probiotics have been reported to exert anticancer effects through various mechanisms, the present study provides correlative observations between *B. bifidum*-derived PFM treatment and changes in miRNA and target gene expression. The precise mechanistic pathway remains hypothetical and requires further functional validation.

In recent years, probiotic microorganisms, particularly *Lactobacillus* and *Bifidobacterium* species have gained considerable attention for their anticancer potential. Their effects are mediated through multiple mechanisms, including the production of antioxidant enzymes, scavenging reactive oxygen species (ROS), heavy metal chelation, and neutralization of carcinogens (17, 18). Additionally, probiotics can regulate cell cycle in cancer cells, suppress proliferation, and enhance sensitivity to apoptosis, all of which are crucial processes in tumor suppression (19).

Colorectal cancer (CRC) is characterized by uncontrolled cell proliferation and evasion of apoptosis, leading to tumor progression. Emerging evidence highlights the capacity of *Bifidobacterium* species to modulate genes involved in apoptotic and anti-apoptotic pathways, suggesting potential therapeutic opportunities (20).

In this study, the post-fermentation medium (PFM) derived from *B. bifidum* exhibited notable cytotoxic effects on HT-29 colorectal cancer cells in a dose-dependent manner with an IC_{50} of approximately 35% (v/v), whereas the bacterial cell extract (CE) showed minimal cytotoxicity. Gene expression and cytotoxic

responses varied across concentrations, likely due to non-linear dose-response effects and differential sensitivities of target genes.

Secreted metabolites present in PFM, such as short-chain fatty acids (SCFAs), proteins, and bacteriocins may play key roles in mediating the observed anticancer activity (21).

Mechanistically, butyrate acts as a histone deacetylase (HDAC) inhibitor, altering chromatin structure and enhancing transcription of pro-apoptotic genes, including BAX and p21, thereby promoting apoptosis and inhibiting cell proliferation (22).

PFM treatment was associated with a moderate but significant upregulation of miR-196a/b-5p. Despite their sequence similarity, miR-196a-5p and miR-196b-5p differ in tissue-specific expression and biological function (6, 23).

Previous studies have reported upregulation of miR-196a in head and neck (23), oral (24), gastric (25), pancreatic (6, 26), cervical (27), and lung (8) cancers, while downregulation of miR-196b in oral (23), gastric (28), colorectal (29), glioblastoma, and leukemia (23) cancers. Both miRNAs are implicated in oncogenic processes, enhancing proliferation, migration, invasion, and resistance to therapy, though tumor-suppressive functions have also been reported in certain contexts, such as melanoma and breast cancer (30). Similarly, chemotherapeutic agents like Epirubicin can induce specific miRNAs, including miRNA-22 and miRNA-331, affecting cell survival and metastasis in triple-negative breast cancer (31).

To evaluate downstream effects, the expression levels of established miR-196 target genes—*HOXB8*, *IGF2BP3*, and *E2F7*—were measured. PFM treatment led to significant downregulation of *HOXB8* and *IGF2BP3*, while *E2F7* remained unchanged. Although these results suggest a potential link between PFM treatment and miR-196a/b-5p-associated regulation of *HOXB8* and *IGF2BP3*, the observations are correlative, and direct causality has not been experimentally confirmed.

The extent of target gene suppression may also be influenced by other bioactive metabolites in PFM, such as SCFAs, and is not proven to be solely mediated by miRNA changes (32). Functional studies using miRNA mimics, inhibitors, or gene knockdown are required to validate these regulatory relationships.

HOXB8 contributes to CRC progression by promoting proliferation and migration via the Wnt/ β -catenin pathway (33). Elevated *HOXB8* correlates with poor prognosis and is observed even in precancerous lesions (34). It promotes epithelial–mesenchymal transition (EMT) and activates STAT3 signaling, enhancing oncogenic drivers like c-Myc and Cyclin D1 (35, 36). Importantly, the ratio of miR-196 to *HOXB8* can modulate sensitivity to FOLFOX4 chemotherapy in CRC, where increased miR-196 levels correlate with reduced *HOXB8* and improved chemotherapeutic response (37).

IGF2BP3, an oncofetal RNA-binding protein, is associated with tumor progression and poor prognosis in multiple cancers (38, 39). It stabilizes oncogenic mRNAs, including *RCC2* and *EGFR*, promoting survival, chemoresistance, and proliferation (40, 41). Bioinformatic analyses identify *IGF2BP3* as a target of miR-196b-5p (42, 43).

This study provides the first evidence that *B. bifidum*-derived PFM downregulates *IGF2BP3* in colorectal cancer cells, potentially via miR-196a/b-5p, though functional validation is necessary.

Conversely, CE treatment did not significantly affect miRNA or target gene expression, reinforcing that secreted metabolites, rather than bacterial cellular components, are likely responsible for the observed anticancer activity. Notably, at 50% PFM, miRNA upregulation and *IGF2BP3* downregulation were slightly less pronounced than at 35%, despite greater cytotoxicity. This non-linear response may result from impaired cellular machinery required for miRNA biogenesis and mRNA regulation under highly cytotoxic conditions.

The present study has several limitations. First, the observed changes in miR-196a/b-5p and their target genes are correlative and do not establish a direct mechanistic link. Second, experiments were

performed using a single colorectal cancer cell line (HT-29), which may limit the generalizability of the findings. Future studies should include functional validation using miRNA mimics, inhibitors, or gene knockdown, as well as testing additional CRC to confirm the selectivity and reproducibility of the observed effects.

The other limitation of the study is the lack of evaluation of PFM and CE effects on normal colorectal epithelial cells. Functional validation of the proposed miRNA-mediated regulatory mechanism was not performed. Future investigations should include non-cancerous colon epithelial cell lines to assess selective toxicity and further clarify safety and therapeutic potential.

5. Conclusion

This study demonstrates that the post-fermentation medium (PFM) derived from *Bifidobacterium bifidum* exerts significant anticancer effects on HT-29 colorectal cancer cells. PFM treatment reduced cell viability in a dose-dependent manner and was associated with upregulation of miR-196a/b-5p and downregulation of their known oncogenic targets, *HOXB8* and *IGF2BP3*. In contrast, the bacterial cell extract (CE) showed negligible effects on these parameters. These findings indicate that fermentation-derived bioactive metabolites may contribute to the observed anticancer activity. However, the observed relationships between PFM, miRNA changes, and target gene expression are correlative and do not establish a direct mechanistic link. The substantial gene downregulation observed may also be influenced by other bioactive compounds in PFM (e.g., SCFAs with HDAC-inhibitory activity), independent of -or only partially dependent on- the modest miRNA changes. Therefore, these mechanistic interpretations remain hypothesis-generating. Future functional studies using miRNA mimics, inhibitors, or gene knockdown are required to validate whether miR-196a/b-5p directly mediates the observed gene expression changes, and *in vivo* investigations are needed before drawing therapeutic conclusions.

6. Declarations

6.1 Acknowledgment

The authors gratefully acknowledge Dr. Shekufe Rezghi Barez (Isfahan University of Medical Sciences, Iran) for her assistance in experimental procedures.

6.2 Ethical Considerations

Not applicable. This study did not involve human participants or animal experiments.

6.3 Authors' Contributions

Somayeh Farahmand: Conceptualization, methodology, software, validation, investigation, data curation, writing — original draft, review & editing, visualization, supervision. Arghavan Sabzevari: Conceptualization, methodology, software, formal analysis, investigation, resources, data curation, writing — original draft, review & editing, visualization, supervision, project administration, funding acquisition. Mohammad Amin Nicknezhad: Conceptualization, review & editing, supervision.

6.4 Conflict of Interests

The authors report there are no competing interests to declare.

6.5 Financial Support and Sponsorship

This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

6.6 Using Artificial Intelligence Tools (AI Tools)

The authors used AI-based language tools (e.g., grammar or style checkers) to improve the readability of the manuscript. All content, interpretations, and conclusions are solely the work of the authors.

References

1. Rawla P, Sunkara T, Barsouk A. Epidemiology of colorectal cancer: incidence, mortality, survival, and risk factors. *Gastroenterol Rev (Prz Gastroenterol)*. 2019;14(2):89-103. [DOI:10.5114/pg.2018.81072] [PMID] [PMCID]
2. Li J, Chu R, Wang C, Li Y, Wu B, Wan J. Microbiome characteristics and *Bifidobacterium longum* in colorectal cancer patients pre-and post-chemotherapy. *Transl Cancer Res*. 2020;9(4):2178. [DOI:10.21037/tcr.2020.03.33] [PMID] [PMCID]
3. Turrone F, Duranti S, Milani C, Lugli GA, van Sinderen D, Ventura M. *Bifidobacterium bifidum*: a key member of the early human gut microbiota. *Microorganisms*. 2019;7(11):544. [DOI:10.3390/microorganisms7110544] [PMID] [PMCID]
4. Turrone F, Milani C, Ventura M, van Sinderen D. The human gut microbiota during the initial stages of life: insights from bifidobacteria. *Curr Opin Biotechnol*. 2022;73:81-7. [DOI:10.1016/j.copbio.2021.07.012] [PMID]
5. Karami P, Goli HR, Abediankenari S, Chandani SR, Jafari N, Ghasemi M, et al. Anti-tumor effects of *Bacteroides fragilis* and *Bifidobacterium bifidum* culture supernatants on mouse breast cancer. *Gene Rep*. 2023;33:101815. [DOI:10.1016/j.genrep.2023.101815]
6. Wu C, Wu Z, Wang L, Chen Y, Huang X, Wang Z, et al. The Modulating Mechanisms of miRNA-196 in Malignancies and Its Prognostic Value: A Systematic Review and Meta-Analysis. *Nutr Cancer*. 2022;74(2):423-36. [PMID] [DOI:10.1080/01635581.2021.1922718]
7. Maralani M, Shanebandi D, Asadi M, Hashemzadeh S, Hajiasgharzadeh K, Mashhadi Abdolahi H, et al. Expression profiles of miR-196, miR-132, miR-146a, and miR-134 in human colorectal cancer tissues in accordance with their clinical significance: Comparison regarding KRAS mutation. *Wien Klin Wochenschr*. 2021;133(21):1162-70. [PMID] [DOI:10.1007/s00508-021-01933-9]
8. Xiong M, Wang P, Pan B, Nie J, Wang S, He B. The diagnostic and prognostic values of microRNA-196a in cancer. *Biosci Rep*. 2021;41(1):BSR20203559. [DOI:10.1042/BSR20203559] [PMID] [PMCID]
9. Maleki M, Golchin A, Javadi S, Khelghati N, Morovat P, Asemi Z, et al. Role of exosomal miRNA in chemotherapy resistance of Colorectal cancer: A systematic review. *Chem Biol Drug Des*. 2023;101(5):1096-112. [DOI:10.1111/cbdd.13947] [PMID]
10. Lederer M, Bley N, Schleifer C, Hüttelmaier S. The role of the oncofetal IGF2 mRNA-binding protein 3 (IGF2BP3) in cancer. In *Seminars in cancer biology 2014 Dec 1 (Vol. 29, pp. 3-12)*. Academic Press. [PMID] [DOI:10.1016/j.semcancer.2014.07.006]
11. Ma Y, Wang S, Bao J, Wang C. Systematic study on expression and prognosis of E2Fs in human colorectal cancer. *Int J Clin Oncol*. 2022;27(2):362-72. [DOI:10.1007/s10147-021-02051-2] [PMID]
12. Elahi Z, Shariati A, Bostanghadiri N, Dadgar-Zankbar L, Razavi S, Norzaee S, et al. Association of *Lactobacillus*, Firmicutes, *Bifidobacterium*,

- Clostridium, and Enterococcus with colorectal cancer in Iranian patients. *Heliyon*. 2023;9(12):e22602. [DOI:10.1016/j.heliyon.2023.e22602] [PMID] [PMCID]
13. Chattopadhyay I, Dhar R, Pethusamy K, Seethy A, Srivastava T, Sah R, et al. Exploring the role of gut microbiome in colon cancer. *Appl Biochem Biotechnol*. 2021;193(6):1780-99. [DOI:10.1007/s12010-021-03498-9] [PMID]
 14. Khosrovan Z, Haghghat S, Mahdavi M. The Probiotic Bacteria Induce Apoptosis in Breast and Colon Cancer Cells: An Immunostimulatory Effect. *Immunoregulation*. 2020;3(1):37-50. [DOI:10.32598/IMMUNOREGULATION.3.1.5]
 15. Nowak A, Zaklos-Szyda M, Rosicka-Kaczmarek J, Motyl I. Anticancer potential of post-fermentation media and cell extracts of probiotic strains: an in vitro study. *Cancers*. 2022;14(7):1853. [PMID] [PMCID] [DOI:10.3390/cancers14071853]
 16. Bustin SA, Benes V, Garson JA, Hellemans J, Huggett J, Kubista M, et al. The MIQE Guidelines: Minimum Information for Publication of Quantitative Real-Time PCR Experiments. 2009. Pp. 611-22. Oxford, United Kingdom: Oxford University Press. [DOI:10.1373/clinchem.2008.112797] [PMID]
 17. Lu K, Dong S, Wu X, Jin R, Chen H. Probiotics in cancer. *Front Oncol*. 2021;11:638148. [DOI:10.3389/fonc.2021.638148] [PMID] [PMCID]
 18. Kaźmierczak-Siedlecka K, Roviello G, Catalano M, Polom K. Gut microbiota modulation in the context of immune-related aspects of *Lactobacillus* spp. and *Bifidobacterium* spp. in gastrointestinal cancers. *Nutrients*. 2021;13(8):2674. [DOI:10.3390/nu13082674] [PMID] [PMCID]
 19. Singh S, Singh M, Gaur S. Probiotics as multifaceted oral vaccines against colon cancer: A review. *Front Immunol*. 2022;13:1002674. [DOI:10.3389/fimmu.2022.1002674] [PMID] [PMCID]
 20. Faghfoori Z, Faghfoori MH, Saber A, Izadi A, Yari Khosroushahi A. Anticancer effects of bifidobacteria on colon cancer cell lines. *Cancer Cell Int*. 2021;21(1):258. [PMID] [PMCID] [DOI:10.1186/s12935-021-01971-3]
 21. Kumar A, Ali A, Kapardar RK, Dar GM, Nimisha, Apurva, et al. Implication of gut microbes and its metabolites in colorectal cancer. *J Cancer Res Clin Oncol*. 2023;149(1):441-65. [PMCID] [DOI:10.1007/s00432-022-04422-2] [PMID]
 22. Saldanha SN, Kala R, Tollefsbol TO. Molecular mechanisms for inhibition of colon cancer cells by combined epigenetic-modulating epigallocatechin gallate and sodium butyrate. *Exp Cell Res*. 2014;324(1):40-53. [PMCID] [DOI:10.1016/j.yexcr.2014.01.024] [PMID]
 23. Dioguardi M, Cantore S, Sovereto D, La Femina L, Caloro GA, Spirito F, et al. Potential role of miR-196a and miR-196b as prognostic biomarkers of survival in head and neck squamous cell carcinoma: a systematic review, meta-analysis and trial sequential analysis. *Life*. 2022;12(8):1269. [DOI:10.3390/life12081269] [PMID] [PMCID]
 24. Cheng AJ, You GR, Lee CJ, Lu YC, Tang SJ, Huang YF, et al. Systemic investigation identifying salivary miR-196b as a promising biomarker for early detection of head-neck cancer and oral precancer lesions. *Diagnostics*. 2021;11(8):1411. [DOI:10.3390/diagnostics11081411] [PMID] [PMCID]
 25. Zheng J, Jiang X, Jiang K, Yan Y, Pan J, Liu F, et al. miR-196a-5p correlates with chronic atrophic gastritis progression to gastric cancer and induces malignant biological behaviors of gastric cancer cells by targeting ACER2. *Mol Biotechnol*. 2023;65(8):1306-17. [DOI:10.1007/s12033-022-00589-8] [PMID]
 26. Prinz C, Fehring L, Frese R. MicroRNAs as indicators of malignancy in pancreatic ductal adenocarcinoma (PDAC) and cystic pancreatic lesions. *Cells*. 2022;11(15):2374. [DOI:10.3390/cells11152374] [PMID] [PMCID]
 27. Chen CF, Hu TQ, Han YX, Zhou XF, Fang QM. Influence of miR-146a and miR-196a2 gene polymorphisms on susceptibility to cervical cancer in patients with high-risk HPV infection. *Chin J Nosocomiol*. 2021;10:1566-70.
 28. Chen L, Tang H, Liu G, Xiao S, Liang D, Ma J, et al. MicroRNA-196b promotes gastric cancer progression by targeting ECRG4. *Anti-Cancer Drugs*. 2021;32(2):127-37. [DOI:10.1097/CAD.0000000000000998] [PMID]
 29. Huang L, Liang D, Zhang Y, Chen X, Chen J, Wen C, et al. METTL3 promotes colorectal cancer metastasis by promoting the maturation of pri-microRNA-196b. *J Cancer Res Clin Oncol*. 2023;149(8):5095-108. [PMID] [PMCID] [DOI:10.1007/s00432-022-04429-9]
 30. Xu C, Gu L. The diagnostic effect of serum miR-196b as biomarker in colorectal cancer.

- Biomed. Rep. 2017;6(1):39-45.
[DOI:10.3892/br.2016.815] [PMID] [PMCID]
31. Farahmand S, SamadiAfshar S, Khalili M, Hosseini RH. Therapeutic implications of Epirubicin-induced miRNA-22 and miRNA-331 upregulation on cell viability and metastatic potential in triple-negative breast cancer. *Hum Gene.* 2025;44:201396.
[DOI:10.1016/j.humgen.2025.201396]
 32. D'Amore T, Zolfanelli C, Lauciello V, Di Ciancia A, Vagliasindi A, Smaoui S, et al. Using postbiotics from functional foods for managing colorectal cancer: Mechanisms, sources, therapeutic potential, and clinical perspectives. *Microorganisms.* 2025;13(6):1335.
[DOI:10.3390/microorganisms13061335] [PMID] [PMCID]
 33. Guo J, Zhang T, Dou D. Knockdown of HOXB8 inhibits tumor growth and metastasis by the inactivation of Wnt/ β -catenin signaling pathway in osteosarcoma. *Eur J Pharmacol.* 2019;854:22-7.
[DOI:10.1016/j.ejphar.2019.04.004] [PMID]
 34. Francis JC, Gardiner JR, Renaud Y, Chauhan R, Weinstein Y, Gomez-Sanchez C, et al. HOX genes promote cell proliferation and are potential therapeutic targets in adrenocortical tumours. *Br J Cancer.* 2021;124(4):805-16.
[DOI:10.1038/s41416-020-01166-z] [PMID] [PMCID]
 35. Jiang S, Wang T, Han Y, Hida T, Afzal MZ, Zhou C, et al. Downregulation of homeobox B8 in attenuating non-small cell lung cancer cell migration and invasion through the epithelial-mesenchymal transition pathway. *Transl Cancer Res.* 2024;13(1):413-22.
[DOI:10.21037/tcr-23-2344] [PMID] [PMCID]
 36. Liu L, Wang L, Li X. The roles of HOXB8 through activating Wnt/ β -catenin and STAT3 signaling pathways in the growth, migration and invasion of ovarian cancer cells. *Cytotechnology.* 2022; 74(1):77-87. [PMID] [PMCID]
[DOI:10.1007/s10616-021-00508-w]
 37. Shen S, Pan J, Lu X, Chi P. Role of miR-196 and its target gene HoxB8 in the development and proliferation of human colorectal cancer and the impact of neoadjuvant chemotherapy with FOLFOX4 on their expression. *Oncol Lett.* 2016; 12(5):4041-7. [DOI:10.3892/ol.2016.5210] [PMID] [PMCID]
 38. Liu H, Zeng Z, Afshar M, Lin C, Wang S, Yang H, et al. Overexpression of IGF2BP3 as a potential oncogene in ovarian clear cell carcinoma. *Front Oncol.* 2020;9:1570.
[DOI:10.3389/fonc.2019.01570] [PMID] [PMCID]
 39. Mancarella C, Scotlandi K. IGF2BP3 from physiology to cancer: novel discoveries, unsolved issues, and future perspectives. *Front Cell Dev Biol.* 2020;7:363. [PMID] [PMCID]
[DOI:10.3389/fcell.2019.00363]
 40. Chen LJ, Liu HY, Xiao ZY, Qiu T, Zhang D, Zhang LJ, et al. IGF2BP3 promotes the progression of colorectal cancer and mediates cetuximab resistance by stabilizing EGFR mRNA in an m6A-dependent manner. *Cell Death Dis.* 2023;14(9): 581. [DOI:10.1038/s41419-023-06099-y] [PMID] [PMCID]
 41. Gou J, Li H, Bi J, Pang X, Li X, Wang Y. Transfer of IGF2BP3 through Ara-C-induced apoptotic bodies promotes survival of recipient cells. *Front Oncol.* 2022;12:801226. [PMID] [PMCID]
[DOI:10.3389/fonc.2022.801226]
 42. Visani M, Marucci G, de Biase D, Giangaspero F, Buttarelli FR, Brandes AA, et al. miR-196B-5P and miR-200B-3P are differentially expressed in medulloblastomas of adults and children. *Diagnostics.* 2020;10(5):265. [PMCID]
[DOI:10.3390/diagnostics10050265] [PMID]
 43. Tang J, Wang S, Weng M, Guo Q, Ren L, He Y, et al. The IGF2BP3-COPS7B axis facilitates mRNA translation to drive colorectal cancer progression. *Cancer Res.* 2023;83(21):3593-610. [DOI:10.1158/0008-5472.CAN-23-0557] [PMID]